



III Subjective probabilities

1. Bayesianism

III.1.1 – The Emperor's New Cloth or The Solution to Everything?



According to a recent book Bayesianism is indeed the “Solution to Everything”



Bayesianism has been of good use to

- Prove that smoking causes lung cancer
- Predict winner in the US presidential elections
- Show that cholesterol causes heart attacks
- Find the author of a 18th century document
- Improve the US insurance industry
- Crack the German Enigma code during WW II
- Track down a lost American H-bomb
- Hunt down Soviet submarines
- Investigate nuclear power safety
- Predict the shuttle Challenger accident
- . . . and much more!


III.1.2 The historical background to Bayesianism



Thomas Simpson
1710-61
Mathematician





David Hume 1711-76
"There is no First Cause"



Richard Price
1723-91
Radical priest

Friend of the
American and
French revolutions

Supported women's
rights



Thomas Bayes
1701-61
Dissident clergyman

LII. *An Essay towards solving a Problem in the Doctrine of Chances.* By the late Rev. Mr. Bayes, F. R. S. communicated by Mr. Price, in a Letter to John Canton, A. M. F. R. S.


1763

Dear Sir,

Read Dec. 23,
1763. **I** Now send you an essay which I have found among the papers of our deceased friend Mr. Bayes, and which, in my opinion, has great merit, and well deserves to be preserved. Experimental philosophy, you will find, is nearly interested in the subject of it; and on this account there seems to be particular reason for thinking that a communication of it to the Royal Society cannot be improper.

The possible roots to Thomas Bayes' 1757 theorem

- How to treat randomness
- Is God the Ultimate Cause?
- How to draw conclusions from observations



The first Bayesian was Simone
de Laplace 1749-1827



Bruno de Finetti
(1906-85)



III.1.3 What is Bayesianism?

Frequentists and Bayesians differ in their definition of “probability”

Classical definition: from tossing coins or dice *Both camps agree*

Frequentist definition: the long-run frequency of a “repeatable event” *Agreement*

Bayesian definition: a person’s degree of belief in an event, given the information available *Sharp disagreements*

Bayesianism consists of two parts, one non-controversial, one *highly controversial*


NON-CONTROVERSIAL: The equation for conditional probabilities

$$\text{prob}(\mathbf{A}|\mathbf{B}) = \frac{\text{prob}(\mathbf{A}) \cdot \text{prob}(\mathbf{B}|\mathbf{A})}{\text{prob}(\mathbf{B})}$$

III.1.4 The non-controversial part – conditional probabilities

Classical probability

10%



0% of cases



3% - “ -




24% - “ -



73% - “ -

60%



22% of cases



43% - “ -



29% - “ -



6% - “ -

Bayesian statistics address the problem:

- What is the unknown proportion of red and white balls?



Draw three balls at random




What conclusions can we
draw about the “true”
probability?

*Bayesian statistics deals with **inverse probabilities***

This can be solved by using
the non-controversial
“Bayes Rule” (to be derived)

$$\text{prob}(\mathbf{A}|\mathbf{B}) = \frac{\text{prob}(\mathbf{A}) \cdot \text{prob}(\mathbf{B}|\mathbf{A})}{\text{prob}(\mathbf{B})}$$

What does $A|B$ mean?




$p(B|A)$ means
probability of B
given A (within the
 A area) = C/A

$p(A|B)$ means
probability of A
given B (within the
 B area) = C/B

III.1.5 Deriving Bayes's equation for single outcomes

Deriving Bayes' theorem



$$p(A) = A/N$$


$$p(B) = B/N$$

$$\frac{p(A)}{p(B)} = \frac{A}{B}$$

$$p(B|A) = C/A$$

$$p(A|B) = C/B$$

Deriving Bayes' theorem



$$A \cdot p(B|A) = C$$

$$B \cdot p(A|B) = C$$

$$p(B) \cdot p(A|B) = p(A) \cdot p(B|A)$$


$$p(A|B) = \frac{p(A) \cdot p(B|A)}{p(B)}$$

III.1.5 The controversial part - subjective probabilities

Bayesianism consists of two parts, one non-controversial, one *highly controversial*

NON-CONTROVERSIAL: The derived equation for conditional probabilities

HIGHLY CONTROVERSIAL: The use of subjective probabilities (degrees of belief) and the use of this term – the “prior”


$$\text{prob}(\mathbf{A}|\mathbf{B}) = \frac{\text{prob}(\mathbf{A}) \cdot \text{prob}(\mathbf{B}|\mathbf{A})}{\text{prob}(\mathbf{B})}$$

The controversy is not about mathematics but about philosophy!

The controversial bit, “the prior” which can even be a **pure guess**

$$p(\mathbf{A}|\mathbf{B}) = \frac{p(\mathbf{A}) \cdot p(\mathbf{B}|\mathbf{A})}{p(\mathbf{B})}$$

The “prior” $p(\mathbf{A})$ + new information $p(\mathbf{B}|\mathbf{A})/p(\mathbf{B})$ updates the **probability of A**, i.e. $p(\mathbf{A}|\mathbf{B})$

Objections to subjective probabilities:

“Science and mathematics are supposed to be objective and not subjective”

The statistical community has since long been divided into Frequentists and Bayesians




Karl Pearson
1857-1936



Ronald S Fisher
1890-1962




Robert O. Schiefer
1914-94



Harold Jeffreys
1891-1989




Jerzy Neyman
1894-1981



Egon Pearson
1895-1980



Leonard J. Savage
1917-71



Howard Raiffa 1924 -




Frequentist approach (the long-established majority):

Probability is defined as the long-run frequency of a “repeatable event”.

It developed a notion of confidence interval, with a probability that it is covering the true value.

Confidence intervals



Bayesian approach

(a growing minority):

also

Probability is defined as a person's degree of belief in an event, given the information available.

It developed a notion of credible interval with a probability that the true value will be within the interval.

Truth

80%

Whereas confidence intervals are based only on the data (**clean**), credible intervals incorporate (**dirty**) problem-specific contextual information from the prior distribution

Frequentists

*Bayes
Rule is
wrong!*

Bayesians

The
Frequentists
have for long
regarded the
Bayesians
as spiteful
heretics

... but it works!

As a consequence three types of Bayesians have developed

- 1. Open Bayesians** (openly using and promoting Bayes theorem)
- 2. Covert Bayesians** (using Bayes' theorem without mentioning it)
- 3. Latent Bayesians** (using Bayes theorem without being aware of it)

You had to “come out” as a Bayesian

III.1.7 Why do we need Bayesianism?

- a) To allow subjective probabilities
- b) To avoid over-confident probabilities
- c) To allow updating of preliminary probabilities

“Far better having an **approximate** answer to the **right** question, which is often vague, than an **exact** answer to the **wrong** question, which can always be made precise”. *John W. Tukey*
(1915-2000)

a1) Updating of subjective probabilities


Denmark-Sweden football

Probability


After 78 minutes:
0 - 1




a2) Updating of subjective probabilities



a3) Updating of subjective probabilities




Intuitive Bayesianism among weather forecasters



Bayesianism is making inroad with NWP modellers



- Combination of Gaussian prior & observation
- Gaussian posterior,
- weights independent of values.



Andrew Lorenc,
2004

b) Updating of preliminary probabilities

Arthur L. Bailey

accountant

had to start some insurance activity in the US in spite of lacking statistics on accidents

Had to start by guessing and then modify them in a Bayesian way



c) Avoids over-confident probabilities

such as Concorde before 2000 being the safest air plane



$$\frac{0}{100\ 000^h} < \frac{1}{1\ 000\ 000^h}$$

Concorde Some other airline

accidents

... after the 2000 crash the most unsafe




$$\frac{1}{100\ 000^h} > \frac{1}{1\ 000\ 000^h}$$

A Bayesian would not have regarded Concorde as the world's safest airplane before 2000

“Laplace Rule of Succession”
prevents us from being overconfident
since p never
takes values
0% or 100%

$$p = \frac{1 + N_{\text{accidents}}}{2 + N}$$



“-Never say never”

Other
suggestions
to “Laplace’s
Rule”

Laplace

$$p = \frac{1 + N_{\text{rain}}}{2 + N}$$

Roulstone
and Smith

$$p = \frac{1/2 + N_{\text{rain}}}{1 + N}$$

Neil Bowler
UKMO

$$p = \frac{1 + N_{\text{rain}}}{3 + N}$$

III:2:8 Controversies over the Bayesian approach

1. Subjective probabilities (unavoidable)
2. “Creditability” intervals (accepted in weather forecasting)
3. Complicated calculations (helped by new computers and the Markov Chain Monte Carlo simulations)

END